

NEW ETHOCUPS™ for easy dosing with accuracy

Introducing orange-flavored ethosuximide
now available in prefilled cups



Ethocups™
(ethosuximide oral solution, USP)

Just open, drink, and throw away.

IMPORTANT SAFETY INFORMATION

What is EthoCups™?

EthoCups™ (ethosuximide oral solution) is a prescription medicine used to treat absence seizures (also called petit mal seizures). It helps reduce seizures by calming abnormal brain activity.

Do not stop taking EthoCups suddenly.

Stopping this medicine without your doctor's advice can lead to serious problems, including more frequent or severe seizures.

Please see additional Important Safety Information throughout this brochure and accompanying full Prescribing Information, including the Medication Guide.

The stress of accurate daily dosing

Making sure your child gets the right dose of their daily absence seizure medication can feel stressful, especially when they're young. It's crucial to get the exact amount correct to manage their condition, and measuring liquid medicine precisely isn't always easy.

More than **40%** In fact, more than 40% of caregivers have made a mistake when dosing their child's liquid medication.

It can also be hard to ensure doses are given accurately during school hours or when away from home—like while traveling or spending time with friends or family. No wonder the **odds of dosing error are nearly 5 times higher** when measuring doses into the cup provided by the medication manufacturer, especially for smaller doses.

NEW ETHOCUPS
makes dosing simple



IMPORTANT SAFETY INFORMATION (CONT'D)

Who should not take EthoCups?

Do not take EthoCups if you are allergic to ethosuximide or other similar seizure medicines (called succinimides).

Serious side effects may include:

1. Suicidal thoughts or behaviors

Seizure medicines may increase the risk of suicidal thoughts or actions.

Call your doctor right away if you or your child has:

- New or worsening depression or mood changes
- Thoughts of suicide or self-harm
- Trouble sleeping or feeling agitated


The ease of ETHOCUPS

The prefilled cup formulation of ETHOCUPS supports precise dosing of ethosuximide.



Reliable dosing

Accurate dosing every time when taken as directed



Simple to use

- Less risk of spills or mess
- Convenient for parents and children



No prep or cleanup

- No measuring or mixing needed
- No repacking or refrigeration



ETHOCUPS has an **orange flavor**, which is popular for kids.

IMPORTANT SAFETY INFORMATION (CONT'D)

Serious side effects may include: (cont'd)

2. Serious blood problems

EthoCups may cause rare but serious blood disorders, which can be life-threatening.

Call your doctor if you notice:

- Fever, sore throat, or frequent infections
- Easy bruising, unusual tiredness, or vaginal bleeding

Your doctor may check your blood regularly during treatment.

Please see additional Important Safety Information throughout this brochure and accompanying full Prescribing Information, including the Medication Guide.

Take ETHOCUPS anytime, anywhere

Directions for ETHOCUPS

Following the initial dose amount prescribed by your or your child's doctor, the amount will gradually increase based on the response to treatment.

ETHOCUPS IS AVAILABLE IN 3 DOSES: 150 mg/3 mL, 250 mg/5 mL, 500 mg/10 mL

Initial Dosing	
3 to 6 years of age	250 mg per day
6 years of age and older	500 mg per day

It's important to take ETHOCUPS exactly as directed by your prescribing doctor. Don't stop taking ETHOCUPS suddenly, change the amount you take, or change other medications without first talking with your doctor.

ETHOCUPS is designed for convenience, so it can easily be taken within any daily routine.



At home



In school



While traveling

IMPORTANT SAFETY INFORMATION (CONT'D)

Serious side effects may include: (cont'd)

3. Severe allergic reactions and skin problems

EthoCups may cause rare but serious reactions such as:

- DRESS, a reaction with rash, fever, and organ problems
- Stevens-Johnson Syndrome (SJS), a severe skin reaction

Seek emergency care if you have:

- Rash, swelling, or blisters
- Yellowing of the skin or eyes
- Sores in your mouth

Safety and side effects

The main ingredient in ETHOCUPS is ethosuximide, a proven treatment for absence seizures that has been prescribed to children with absence seizures for more than 50 years.

When taking ETHOCUPS, it's common to experience mild side effects, which may include:

- Headache
- Fatigue
- Irritability
- Hiccups
- Nausea
- Vomiting
- Abdominal discomfort
- Weight loss or anorexia

Less frequent but more serious side effects include psychiatric disturbances, dermatologic conditions, blood conditions, abnormal liver or kidney function, bleeding, or nearsightedness.

It's important to stop taking ETHOCUPS and call your doctor immediately at the first sign of a rash, unless it is clearly not drug related.

For information on other possible side effects, read the Important Safety Information in this brochure.

IMPORTANT SAFETY INFORMATION (CONT'D)

Serious side effects may include: (cont'd)

4. Liver and kidney problems

Your doctor may test your liver and kidney function during treatment.

Call your doctor if you feel very tired, develop dark urine, or experience swelling in the arms or legs.

Please see additional Important Safety Information throughout this brochure and accompanying full Prescribing Information, including the Medication Guide.

IMPORTANT SAFETY INFORMATION

What is EthoCups™?

EthoCups™ (ethosuximide oral solution) is a prescription medicine used to treat absence seizures (also called petit mal seizures). It helps reduce seizures by calming abnormal brain activity.

Do not stop taking EthoCups suddenly.

Stopping this medicine without your doctor's advice can lead to serious problems, including more frequent or severe seizures.

Who should not take EthoCups?

Do not take EthoCups if you are allergic to ethosuximide or other similar seizure medicines (called succinimides).

Serious side effects may include:

1. Suicidal thoughts or behaviors

Seizure medicines may increase the risk of suicidal thoughts or actions.

Call your doctor right away if you or your child has:

- New or worsening depression or mood changes
- Thoughts of suicide or self-harm
- Trouble sleeping or feeling agitated

2. Serious blood problems

EthoCups may cause rare but serious blood disorders, which can be life-threatening.

Call your doctor if you notice:

- Fever, sore throat, or frequent infections
- Easy bruising, unusual tiredness, or vaginal bleeding

Your doctor may check your blood regularly during treatment.

3. Severe allergic reactions and skin problems

EthoCups may cause rare but serious reactions such as:

- DRESS, a reaction with rash, fever, and organ problems
- Stevens-Johnson Syndrome (SJS), a severe skin reaction

Seek emergency care if you have:

- Rash, swelling, or blisters
- Yellowing of the skin or eyes
- Sores in your mouth

4. Liver and kidney problems

Your doctor may test your liver and kidney function during treatment.

Call your doctor if you feel very tired, develop dark urine, or experience swelling in the arms or legs.

5. Mood or behavior changes

EthoCups may cause:

- Irritability, hallucinations, or unusual thoughts
- Trouble concentrating, night terrors, or sleep changes
- Aggression or worsening depression

6. Other seizure types may worsen

In some people, EthoCups may make other types of seizures worse. Tell your doctor if your seizures change.

7. Lupus (autoimmune reaction)

Rare cases of lupus have occurred. Your doctor may monitor for symptoms such as joint pain, rash, or fatigue.

Common side effects include:

- Nausea, vomiting, or stomach pain
- Loss of appetite or weight loss
- Diarrhea
- Hiccups
- Sleepiness, dizziness, or feeling tired
- Headache or problems with coordination

Before taking EthoCups, tell your doctor if you:

- Have liver or kidney problems
- Have or have had mood or mental health conditions
- Are pregnant or plan to become pregnant
- Are breastfeeding

Use in children

EthoCups is not recommended for children under 3 years old.

Driving and using machines

EthoCups may cause drowsiness or dizziness. Do not drive or operate heavy machinery until you know how the medicine affects you.

How should I take EthoCups?

- Take exactly as your doctor prescribes
- Don't skip doses or stop suddenly
- If you miss a dose, take it as soon as you remember unless it's almost time for your next dose
- Your doctor may adjust your dose over time

How should I store EthoCups?

- Store at room temperature (68°F to 77°F)
- Keep away from light and freezing
- Keep tightly closed and out of reach of children

Reporting side effects

Call your doctor for medical advice about side effects.

You may also report side effects to the FDA at **1-800-FDA-1088** or visit www.fda.gov/medwatch.

Please see the full Prescribing Information, including the Medication Guide.

Just open, drink, and throw away.

Easy, accurate dosing with strong support



Specialty pharmacy support
Dedicated support staff to help
with insurance and refills, plus
free home delivery



Manufactured in the USA
Consistent, high-quality product
without shipment delays



Financial support

\$0

Patients with commercial
insurance coverage pay
as little as \$0

\$15

Patients without
commercial insurance pay
no more than \$15

**If your child is taking ethosuximide or oral solutions for
absence seizures, ask their doctor about ETHOCUPS.**

IMPORTANT SAFETY INFORMATION (CONT'D)

Serious side effects may include: (cont'd)

6. Other seizure types may worsen

In some people, EthoCups may make other types of seizures worse. Tell your doctor if your seizures change.

**Please see additional Important Safety Information throughout this brochure and
accompanying full Prescribing Information, including the Medication Guide.**